

Early View Article: Online published version of an accepted article before publication in the final form.

Journal Name: Edorium Journal of Microbiology

Type of Article: Original Article

Title: Prevalence and Intensity of Urinary Schistosomiasis and their Effects on Packed Cell Volume of Pupils in Jaba LGA, Kaduna State, Nigeria

Authors: Henry Gabriel Bishop, Helen Ileigo Inabo, Elijah Ekah Ella

doi: To be assigned

Early view version published: October 3, 2016

**How to cite the article:** Bishop HG, Inabo HI, Ella EE. Prevalence and Intensity of Urinary Schistosomiasis and their Effects on Packed Cell Volume of Pupils in Jaba LGA, Kaduna State, Nigeria. Edorium Journal of Microbiology. Forthcoming 2016.

Disclaimer: This manuscript has been accepted for publication. This is a pdf file of the Early View Article. The Early View Article is an online published version of an accepted article before publication in the final form. The proof of this manuscript will be sent to the authors for corrections after which this manuscript will undergo content check, copyediting/proofreading and content formatting to conform to journal's requirements. Please note that during the above publication processes errors in content or presentation may be discovered which will be rectified during manuscript processing. These errors may affect the contents of this manuscript and final published version of this manuscript may be extensively different in content and layout than this Early View Article.

1 **TYPE OF ARTICLE:** Original Article

2

3 **TITLE:** Prevalence and Intensity of Urinary Schistosomiasis and their Effects on  
4 Packed Cell Volume of Pupils in Jaba LGA, Kaduna State, Nigeria

5

6 **AUTHORS**

7 Henry Gabriel Bishop<sup>1</sup>, Helen Ileigo Inabo<sup>2</sup>, Elijah Ekah Ella<sup>3</sup>

8

9 **AFFILIATIONS:**

10 <sup>1</sup>BSc, MSc. (in view), Department of Microbiology, Faculty of Science, Ahmadu Bello  
11 University, Zaria, Kaduna State, Nigeria, gabrielhenrybishop@gmail.com

12 <sup>2</sup>BSc, MSc., PhD., Department of Microbiology, Faculty of Science, Ahmadu Bello  
13 University, Zaria, Kaduna State, Nigeria, heleninabo@yahoo.co.uk

14 <sup>3</sup>BSc, MSc., PhD., Department of Microbiology, Faculty of Science, Ahmadu Bello  
15 University, Zaria, Kaduna State, Nigeria, elijahella@yahoo.com

16

17 **CORRESPONDING AUTHOR DETAILS**

18 Henry Gabriel Bishop

19 Department of Microbiology, Faculty of Science, Ahmadu Bello University, Zaria,  
20 Kaduna State, Nigeria, 810001

21 Email: gabrielhenrybishop@gmail.com

22

23 **Short Running Title:** Effects of Urinary Schistosomiasis on Packed Cell Volume of  
24 Pupils

25

26 **Guarantor of Submission:** The corresponding author is the guarantor of  
27 submission

28

29

30

31

32 **TITLE:** Prevalence and Intensity of Urinary Schistosomiasis and their Effects on  
33 Packed Cell Volume of Pupils in Jaba LGA, Kaduna State, Nigeria

34

## 35 **ABSTRACT**

36

### 37 **Aims**

38 Urinary schistosomiasis is a persistent health burden among African children. They  
39 are mostly unaware of the risks of transmission of schistosomiasis via cercariae-  
40 infested water bodies and hence more infections occur. This study was aimed at  
41 assessing the prevalence and intensity of urinary schistosomiasis and their effects  
42 on packed cell volume (PCV), and the association of the disease with some socio-  
43 demographic and risks factors among pupils in Jaba LGA of Kaduna State, Nigeria.

44

### 45 **Methods**

46 Awareness lectures were organized in pre-selected public primary schools. A total of  
47 505 pupils volunteered to participate in the study. From each volunteered pupil, 10ml  
48 urine and 2ml blood samples were collected. The urine samples were concentrated  
49 by centrifugation; the sediments were examined microscopically using 10x and 40x  
50 objectives for *Schistosoma haematobium* egg(s) while count/10ml urine was  
51 recorded. Intensity categories were taken as light infection (with <50eggs/10ml urine)  
52 and heavy infection (with >50eggs/10ml urine). Blood samples were used for PCV  
53 determination by microhaematocrit centrifuge technique (HCT); anaemic PCV was  
54 <34%, normal PCV was  $\geq 34 \leq 45\%$ , high PCV was  $\geq 46\%$ . Results and data on socio-  
55 demographic and risk factors were subjected to various statistical analyses at  $p=0.05$   
56 with IBM SPSS Version 21.

57

### 58 **Results**

59 An overall prevalence of 12.3% was obtained for urinary schistosomiasis. Three  
60 villages (Bitaro, Ankun and Kwoi) recorded the highest prevalence of the infection.  
61 However, the infection was absent in two villages (Nok and Sambang). The highest  
62 intensity among the pupils was 204 eggs/10ml urine. The Central Area had the  
63 highest mean intensity of 6.77 eggs/10ml urine. Areas of highest prevalence did not

64 coincide with areas of highest intensity. The infection and its intensity were higher  
65 among the females (15.5%, 4.18 eggs/10ml urine) than the males (9.1%, 1.22  
66 eggs/10ml urine) respectively. Similarly, the females had higher light and heavy  
67 infections than the males. There was an observed increase of urinary  
68 schistosomiasis with increase in pupils' class. Both the infection and its intensity had  
69 gradual 'wave-like' increases with rise in age of the pupils. Only two signs/symptoms  
70 (painful micturition, urine colour), and one risk factor ('Fadama' farming) were  
71 statistically associated with urinary schistosomiasis. The prevalence of anaemia was  
72 found to be 8.1% while 37.6% of the pupils had normal PCV; the remaining study  
73 population had abnormally high PCV. There was a statistically significant  
74 association between urinary schistosomiasis and anaemia among the pupils ( $\chi^2 =$   
75 11.870;  $df = 2$ ;  $p = 0.003$ ). Though anaemia was recorded both among the infected  
76 and uninfected pupils, a higher occurrence of the anaemia (17.7%) was observed in  
77 pupils infected with urinary schistosomiasis than those who were not infected (6.8%).  
78 The cause of the anaemia in the later may be due to other diseases. There was  
79 higher level of non-occurrence of urinary schistosomiasis in pupils with high PCV  
80 (56.4%). Heavy infections with urinary schistosomiasis among the pupils, with a  
81 statistical significance ( $\chi^2 = 12.807$ ;  $df = 4$ ;  $p = 0.012$ ) led to higher occurrence of  
82 anaemia of 20.0% than light infections which caused 17.2% of anaemia.

83

#### 84 **Conclusion**

85 With an overall prevalence of 12.3% and varying levels of intensity, urinary  
86 schistosomiasis is still prevalent in Nigeria which calls for concerted efforts to  
87 eradicate its menace in all affected regions. Whatever that affects the health of  
88 children should not be neglected. The female pupils were significantly more affected  
89 than the male pupils and hence are predisposed to further complications like female  
90 genital schistosomiasis (FGS) and bladder cancers. The disease is associated with  
91 painful micturition and red/brown-coloured urine. Farming on 'Fadama' (i.e.,  
92 waterlogged) farms enhances the acquisition of the disease. Heavy infection with the  
93 worms exacerbates the anaemia in children. There was a total unawareness of the  
94 disease in Jaba LGA of Kaduna State, Nigeria, which is a major promoter of  
95 exposure to the cercariae of the schistosomes during water-contact activities.

96 **Keywords:** Schistosoma haematobium, urine, unawareness, intensity, pupils, Jaba  
97 LGA, Nigeria

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

EARLY VIEW

127 **TITLE:** Prevalence and Intensity of Urinary Schistosomiasis and their Effects on  
128 Packed Cell Volume of Pupils in Jaba LGA, Kaduna State, Nigeria

129

### 130 **INTRODUCTION**

131 Schistosomiasis is a chronic parasitic disease caused by blood flukes [1], [2]. These  
132 blood flukes (or trematode worms) are prime examples of complex multicellular  
133 pathogens that flourish in human hosts despite the presence of immune responses  
134 mounted against them [3]. Adult schistosomes possess a range of adaptations  
135 which enable them to partially overcome human host's defenses and possibly live  
136 and reproduce for many years in the host who becomes relatively resistant to new  
137 infections [4]. These worms have complex and indirect life cycles involving  
138 intermediate (snail) hosts and definitive hosts.

139 Majorly, populations of tropical and subtropical countries, especially children who  
140 indulge in water-based activities in unsafe or cercarial-infested water bodies are  
141 affected [2, 5, 6-8]. The three main species of *Schistosoma* that cause diseases in  
142 man are: *Schistosoma mansoni*, *Schistosoma haematobium* (*S. haematobium*) and  
143 *Schistosoma japonicum* [9, 10]. Other distributions of *Schistosoma intercalatum* in  
144 Central Africa and *Schistosoma mekongi* in Cambodia and Lao People's Democratic  
145 Republic can also cause human infections [11, 12]. Among all the species of  
146 schistosomes, *S. haematobium* is the only cause of urinary schistosomiasis; all the  
147 others cause intestinal schistosomiasis [6, 10].

148 Urinary schistosomiasis is diagnosed by microscopic detection of *S. haematobium*  
149 eggs in urine as a gold standard [6], [13]. Humans become infected by penetration  
150 (or dermo-invasion) of intact skin by active cercariae which are attracted to the  
151 warmth of body and skin lipids [14].

152 *Schistosoma haematobium* has been regarded as a 'neglected schistosome' [15-17]  
153 despite its implication in HIV/AIDS co-infection and a burden of bladder cancer  
154 development. Though most of the infected individuals in endemic areas of Nigeria  
155 suffer from light infections, the disease adversely impacts on the economic and  
156 general health conditions of the affected communities [18, 7]. Consequently, the  
157 workforce is affected due to weakness and lethargy, and the academic performances  
158 of school children are affected [7], 19, 20].

159 **MATERIALS AND METHODS**

160

161 **Materials and methods**

162

163 **Study area**

164 The study was conducted in some selected and consented primary schools in Jaba  
165 Local Government Area (LGA) of Kaduna State, Nigeria. The area is located in the  
166 Northern hemisphere on Latitude 9° 19' 47"N to Latitude 9° 36' 35"N, and in the East  
167 on Longitude 7° 56' 24"E to Longitude 8° 12' 36"E. The area is occupied by the Ham  
168 People, a people notable for the rich Nok culture that possesses the Nok Terracotta  
169 carbon-dated to about 2000-2500 years ago. 'Kwain' (or Kwoi which means  
170 'Community of the United') serves as the political capital of the LGA. It has many  
171 villages that include Bitaro, Nok, Kwoi, Zshiek (Kurmin Musa), Dung (also called  
172 Jaban Kogo), Chori, Fai, Ketere, Sambang Gida, Sambang Daji, Dura, Ankun, Gora,  
173 Kurmi Danagana, Tunga and many other Ham settlements in the Southern parts of  
174 Kaduna State [21]. The people of the area are predominantly farmers; they cultivate  
175 large quantities of gingers, *Digitaria exilis*, (popularly called 'acha' or 'hungry rice'),  
176 cocoyam, guinea corn, millet and maize among many others.

177

178 **Awareness talks and enrollment of volunteers**

179 Awareness talks on schistosomiasis, its danger, transmission, control and prevention  
180 were delivered in pre-selected/consented primary schools in Jaba LGA. The pupils  
181 were informed of the need to volunteer to be part of this study by willingly submitting  
182 their urine and blood samples for laboratory diagnoses. Those pupils that  
183 volunteered to be part of the study were 505 in number; they were given consent  
184 forms to present to their parents/guardians for full permission to enroll them.  
185 Confidentiality was applied on all data collected from them and result of laboratory  
186 tests was issued to each participated pupil. Those that had infection(s) were  
187 immediately referred to the hospital for medical attention.

188

189

190

**191 Urine and blood samples collection**

192 The pupils who consented to participate in the study were briefed and guided on how  
193 to collect 10ml of their urine into provided sterile (wide-mouth) sampling bottles with  
194 screw caps between 10am -1pm [22]. Then 2ml of venous blood were collected into  
195 5ml EDTA K-3 bottle using new sterile syringe and needle for each volunteered pupil  
196 that had submitted a urine sample. The urine samples were screened away from  
197 sunlight by enclosing them in dark polythene bags. Both sample types from each  
198 pupil were simultaneously labelled and placed into separate ice containers. The  
199 samples were taken for analysis at the Bacteriology/Parasitology Laboratory in the  
200 Department of Microbiology, Faculty of Science, Ahmadu Bello University Zaria,  
201 Nigeria.

202

**203 Structured questionnaires administration**

204 Those pupils who submitted urine and blood samples were administered structured  
205 questionnaires. The questionnaires captured some socio-demographic and risk  
206 factors associated with urinary schistosomiasis. Assistance by respective class  
207 teachers and head teachers were sought for interpretation from English language  
208 into easily understood dialect of the study area.

209

**210 Method for determination of packed cell volume (PCV)**

211 The blood samples were brought out of the cold container and allowed to reach the  
212 room temperature upon arrival at the laboratory. The PCV of each pupil was  
213 determined by the microhaematocrit centrifuge technique (HCT). Two plain capillary  
214 tubes were filled with a blood sample to three-fourth their heights and sealed  
215 carefully by means of Bunsen flame to the (2 mm) red demarcation on each tube.  
216 The tubes were spun in the microhaematocrit centrifuge at relative centrifugation  
217 force (RCF) of 12,000-15,000 xg for 5 minutes, after which the PCV were read by  
218 correctly adjusting the red-packed-cells columns on the Haematocrit Reader and an  
219 average of the two values recorded [22]. Anaemic PCV was <34%, normal PCV was  
220  $\geq 34 \leq 45\%$ , high PCV was  $\geq 46\%$  [22].

221

222



**223 Detection and quantification of Schistosoma haematobium eggs in urine**

224 Each urine sample was gently shaken to stir up any sediment in the sampling bottle  
225 and transferred into a labelled centrifuged tube. Normal saline was used to rinse the  
226 bottle where sediments still remained. Centrifugation was performed at a speed of  
227 3000rpm for 3-5 minutes and the supernatant was carefully decanted [22]. The  
228 sediments obtained in the centrifuge tube was tapped on the bench and mixed by  
229 gentle shaking. A Pasteur pipette was used to transfer all the sediments unto a  
230 clean, grease-free glass slide. A drop of Lugol's iodine solution was added and a  
231 cover slip was placed over the wet mount and positioned under the light microscope.  
232 The entire wet mount was screened for egg(s) of Schistosoma haematobium using  
233 10x and 40x objectives, while count of eggs/10ml of urine was taken [22]. Where the  
234 sediments from a sample could not be contained in a single wet mount, multiple wet  
235 mounts were made from such a sample and the egg counts pooled together.

236

**237 Statistical analysis**

238 Data collected of socio-demographic and risk factors of urinary schistosomiasis  
239 together with laboratory findings were subjected to Chi Square ( $\chi^2$ ) and Likelihood  
240 ratio (LR) analyses with the IBM SPSS Statistics Version 21 at  $p=0.05$ . Final results  
241 were presented in charts and tables.

242

**243 RESULTS**

244 Out of 505 urine samples of pupils examined, Schistosoma haematobium ova were  
245 detected in 62 of the samples with a prevalence of 12.3 % (Figure 1). The eggs of  
246 this parasite were yellow-brown, oval in shape with terminal spines; some of the  
247 eggs were shorter but majority were slender (Plate I).

248 The level of intensity of urinary schistosomiasis was determined by obtaining counts  
249 of eggs/10 ml of urine sample and their simple average calculated as shown in Table  
250 1. The highest mean intensity was found among pupils in Primary 6, followed by  
251 those in Primary 4 and Primary 2 respectively. The female pupils in all the classes  
252 had higher intensity than the corresponding male pupils. The highest mean intensity  
253 in the females was 9.74 eggs/10 ml urine which occurred among the female pupils in  
254 Primary 6; whereas that of the males was 2.91 eggs/10 ml urine and it occurred

255 among the male pupils in Primary 4. There was no statistical significance ( $\chi^2 = 6.850$ ;  
256  $df = 5$ ;  $p = 0.232$ ) in the class distribution of the disease, but higher occurrence was  
257 obtained in the senior class category (Primary Class 4-6) and lower in the junior  
258 class category (Primary Class 1-3).

259 Gender was a statistically significant factor in the occurrence of urinary  
260 schistosomiasis ( $\chi^2 = 4.926$ ;  $df = 1$ ;  $p = 0.026$ ) as indicated in Table 2. There was a  
261 remarkable higher occurrence of the infection among the female pupils (15.5%) than  
262 in the male pupils (9.1%). Hence, female pupils were about twice more infected with  
263 urinary schistosomiasis than the male pupils.

264 This study found that in an independent t-test pupils within the age of  $10.96 \pm 2.15$   
265 years had statistically significant occurrence of urinary schistosomiasis compared to  
266 pupils of  $9.70 \pm 2.17$  years ( $t = 4.251$ ,  $df = 503$ ,  $p = 0.000$ ). The occurrence of urinary  
267 schistosomiasis increased in a 'wave-like' fashion with rise the pupils' age. Though  
268 the highest occurrence of 62.1% was found among the older pupils of 15 years of  
269 age, no occurrence was recorded in pupils within 4-6 years age bracket (Figure 2).

270 The mean of counts of eggs/10ml urine adopted a 'wave-like' pattern of increase. No  
271 egg was detected in urine samples of pupils within 4-6 years age bracket. The mean  
272 of counts peaked at age 15 years (Figure 3), which was similar to the age  
273 prevalence pattern in Figure 2. However, the Spearman's correlation ( $r_s$ ) and  
274 Pearson product-moment correlation ( $r$ ) run to determine the relationship between  
275 age of pupils and *Schistosoma haematobium* egg count/10ml of urine (i.e., intensity)  
276 in Jaba LGA showed weak, positive correlations, which were statistically significant  
277 ( $r_s = 0.172$ ,  $P = 0.000$ ;  $r = 0.131$ ,  $P = 0.003$ ).

278 The prevalence of urinary schistosomiasis according to sampling locations in Jaba  
279 LGA showed high statistical significance ( $\chi^2 = 38.599$ ;  $df = 9$ ;  $p = 0.000$ ). The highest  
280 occurrence was found among pupils from Bitaro (23.2%) followed by Ankun (22.2%)  
281 and Kwoi (20.3%), whereas no occurrences were recorded among pupils from Nok  
282 (0.0%) and Sambang (0.0%) as shown in Table 3.

283 The mean intensity of *Schistosoma haematobium* eggs in urine samples of pupils  
284 from the 10 different sampling locations were analysed by means of ANOVA.  
285 Although no statistical significance was obtained (since  $F = 1.419$ ;  $df = 9$ ;  $p = 0.177$ ),

286 the highest means of counts of eggs/10ml of urine were 6.77 and 4.88 among pupils  
287 from the Central Area and Bitaro respectively (Table 3).

288 The level of urinary schistosomiasis among pupils versus sampling locations showed  
289 statistical significance ( $\chi^2 = 50.094$ ,  $df = 18$ ,  $p = 0.000$ ). This was categorized as  
290 'light' and 'heavy' infections, with egg counts of  $<50$ eggs/10ml and  $\geq 50$ eggs/10ml  
291 urine respectively. Light infections occurred most among the pupils from Bitaro  
292 (44.2%), followed by those from Kwoi (21.2%) and Ankun (9.6%). Heavy infections  
293 were most found among pupils from the Central Area (40.0%), followed by those  
294 from Bitaro (30.0%). Light infection was absent among pupils from Nok and  
295 Sambang. Also, heavy infection was not found in pupils from Chori, Dura, Gora, Nok  
296 and Sambang (Table 4).

297 Table 5 shows the age-related distribution of level of urinary schistosomiasis among  
298 the pupils in Jaba LGA (with  $\chi^2 = 44.715$   $df = 22$ ;  $p = 0.003$ ). Both heavy and light  
299 infections occurred mostly in pupils of age 15 years. There was no heavy infection  
300 among pupils in age 4, 5, 6, 8, 11 and 14 years. Also, only pupils in age 4, 5 and 6  
301 did not have light infections.

302 The prevalence of anaemia among pupils in Jaba LGA was found to be 8.1%, while  
303 37.6% of the pupils had normal PCV. The remaining study population (54.3%) had  
304 abnormally high PCV (Figure 4).

305 There was a statistically significant association between urinary schistosomiasis and  
306 anaemia among the pupils ( $\chi^2 = 11.870$ ;  $df = 2$ ;  $p = 0.003$ ). Though anaemia was  
307 recorded both among those with urinary schistosomiasis and uninfected pupils, a  
308 higher occurrence of the anaemia (17.7%) was observed in pupils infected with  
309 urinary schistosomiasis than those who were not infected (6.8%). The cause of the  
310 anaemia in the later may be due to other diseases as indicated in Table 6.

311 Heavy infections with urinary schistosomiasis among the pupils, with a statistical  
312 significance ( $\chi^2 = 12.807$ ;  $df = 4$ ;  $p = 0.012$ ) led to higher occurrence of anaemia of  
313 20.0% than light infections which caused 17.2% of anaemia as indicated in Table 7.

314 The relationship between the level of urinary schistosomiasis and the gender of the  
315 pupils in Jaba LGA was insignificant ( $\chi^2 = 5.166$ ;  $df = 2$ ;  $p = 0.076$ ), but female pupils  
316 had higher proportions of light and heavy infections than the male pupils (Table 8).

317 The level of the urinary schistosomiasis did not influence the onset of (visible)  
318 haematuria among the pupils in this study ( $\chi^2 = 3.105$ ;  $df = 2$ ;  $p = 0.212$ ). All the  
319 pupils with heavy infections did not present with haematuria. Generally, 4.5% of the  
320 pupils presented with haematuria but no eggs of *Schistosoma haematobium* were  
321 recovered; the haematuria may be due to parasitic or other non-parasitic origins.  
322 However, 9.6% of the pupils with light infections presented with haematuria (Table  
323 9).

324 In Table 10, five risk factors of urinary schistosomiasis among school pupils were  
325 considered in this study. There was a statistically important association between  
326 urinary schistosomiasis and 'Fadama' farming ( $\chi^2 = 14.300$ ;  $df = 1$ ;  $p = 0.000$ ). The  
327 pupils that made up the study population were unaware of urinary schistosomiasis.  
328 Pupils that wash their clothes in rivers/streams had more infections (14.7%) than  
329 those who wash their clothes at home (10.8%).

330 Appearances of some of the rivers in Jaba LGA of Kaduna State in Nigeria are  
331 shown in Plate II. Most of the rivers are surrounded by immediate wet fields locally  
332 referred as 'Fadama' for rice, cocoyam and sugarcane farming. The rivers are used  
333 for swimming, fishing, irrigation and washing by pupils and the members of the  
334 communities. The photograph was taken in the late evening when activities were not  
335 on-going.

336 The level of formal education of the pupils' parents had no statistical link with  
337 acquisition of urinary schistosomiasis by their children/wards ( $P > 0.05$ ). However,  
338 lowest detection of *Schistosoma haematobium* eggs was observed in pupils whose  
339 parents did not acquire any formal education. Occurrence of the infection among the  
340 pupils increased as fathers' level of formal education increased (Table 11).

341 From data obtained via administered structured questionnaires on signs and  
342 symptoms of urinary schistosomiasis, only painful micturition ( $\chi^2 = 5.135$ ;  $df = 1$ ;  $p =$   
343  $0.023$ ) and cloudy brown/red-colored urine ( $\chi^2 = 20.604$ ;  $df = 4$ ;  $p = 0.000$ ) had  
344 statistical relevance with the disease as shown in Table 12. Even though pupils with  
345 fever had more infection with *Schistosoma haematobium* (22.2%) than those without  
346 fever, it was insignificant together with the presence of abdominal pain and frequent  
347 urination.

348

349 **DISCUSSION**

350 In this study, the presence of urinary schistosomiasis was established by the  
351 detection of terminal spine eggs in urine samples of the pupils. Cheesbrough [22]  
352 had stated that diagnosis of schistosomiasis is by detection of eggs in urine or stool,  
353 which remains the gold standard.

354 The prevalence of 12.3% for urinary schistosomiasis among the pupils was much  
355 lower than many other findings which had claimed an endemicity of schistosomiasis  
356 in Nigeria. The prevalence in this study was lower than 31.3% reported in Abuja-  
357 Nigeria [23]. Other prevalence reports include 79.4% among children of Ezza-North  
358 LGA of Ebonyi, Nigeria [20], 41.5% in Benue-Nigeria [24], 78.4% in Lagos-Nigeria  
359 [25], 34.1% in Enugu-Nigeria [26], 48.7% in Borno-Nigeria [27], and 19.5% in  
360 Kaduna-Nigeria [28]. Compared to these high-prevalence areas, Jaba LGA had a  
361 lower burden of urinary schistosomiasis. This could be due to the fact that the area  
362 has relatively fewer fast-flowing streams and rivers that may not provide adequate  
363 breeding zone for cercariae and snails (especially *Bulinus* spp and *Physopsis* spp).  
364 Also many of the streams/rivers in the LGA are used for bathing and washing of  
365 clothes by many members of the communities. Detergents and soaps have adverse  
366 effects on cercariae/snails; hence these agents help in shortening the lifecycle and  
367 propagation of cercariae [29].

368 Also, this study uncovered a lesser prevalence of urinary schistosomiasis as  
369 compared to 15.2% prevalence reported in Mali by [30]. A prevalence of 20.7% was  
370 found in a retrospective study in Kumasi [31] and 30.7% had also been reported in  
371 Ghana [32]. A considerably higher prevalence of 35.9% was found in Ethiopia [33],  
372 and a prevalence of 42% among primary school children in South Africa [34].  
373 Although a relatively low prevalence was obtained in this study, it is still higher than  
374 reported 4.5% prevalence in Abini and the absence of the infection in Ukwelo-Obudu  
375 communities of Cross River-Nigeria [35], and reported 6.3% prevalence among  
376 children of Ngbo-West LGA in Ebonyi-Nigeria [20], and 8.3% in Kano-Nigeria [36].

377 The statistically significant higher prevalence (of 15.5%) of urinary schistosomiasis  
378 amongst the females than in the males (9.1%) in this study is among a few reports of  
379 higher occurrence of schistosomiasis amongst the females. This finding supports the  
380 work of Oluwasogo and Fagbemi [25], in which females (60.9%) were more infected

381 than males (39.2%). But this observed phenomenon opposes many findings which  
382 had suggested higher risks on the males [20, 23, [26, 28, 37]. Most of those studies  
383 that had implicated the male gender with higher risk of *Schistosoma* infections had  
384 been conducted in locations where the females were restricted from swimming  
385 activity. In Jaba LGA, both the females and the males had no traditional/religious  
386 restrictions from participating in swimming activity. In addition, there was an unequal  
387 exposure risks between the genders; females participated more in rice farming and  
388 other 'Fadama' activities. Individuals with heavy infections are most prone to chronic  
389 complications [38]. By extension, the females are not only at a higher risk of  
390 developing complications like female genital schistosomiasis (FGS), but they stand  
391 the risks of decreased fertility, abortions, vaginal discharge, contact bleeding and  
392 increased risk of HIV infection in concomitant urinary schistosomiasis [39].

393 Many reports on the prevalence of schistosomiasis had focused on age-group  
394 prevalence. This type of disease-reporting has not explained the occurrences in  
395 each age. Furthermore, conclusions made on an age-group may not adequately  
396 define a particular age characteristics. Hence, in this study the prevalence of urinary  
397 schistosomiasis for each age was calculated, which gave a 'wave-like' pattern of  
398 elevation with increase in ages of the pupils. The absence of the infection in pupils  
399 within ages 4, 5 and 6 years (unlike the older pupils) could be due to the fact they  
400 are too young to engage in constant water-based activities like swimming, wading  
401 or 'Fadama' farming. Older pupils can engage freely and/or indiscriminately in water  
402 activities [37].

403 Aside the obviously high prevalence of urinary schistosomiasis in the females, this  
404 study revealed a higher percentage of heavy infections among the females (70.0%)  
405 than among the males (30.0%). The highest mean intensity of 9.74 eggs/10 ml urine  
406 among the females occurred among pupils in Primary 6; however, in the males it  
407 was 2.91 eggs/10 ml urine found among pupils in Primary 2. In an overall, the  
408 females had higher mean intensity (of 4.18 eggs/10 ml urine) than males (1.22  
409 eggs/10 ml urine). An inclusion of behavioral difference between the genders could  
410 be used to explain reasons for the higher intensity/level of urinary schistosomiasis  
411 among the female pupils. Schistosomes are attracted to skin lipids [14] and since  
412 females (including those that are pupils) often apply high-scented perfumes, body

413 creams and oils, they stand a higher chance of been repeatedly infected during  
414 water-contact activities. Though Bigwan et al. [37] had linked a higher prevalence of  
415 schistosomiasis in males to some socio-cultural practices such as bathing, washing,  
416 swimming, irrigation-farming and fishing in rivers/ponds, the reverse was the case in  
417 this study. Repeated infection episodes enhance the development of heavy  
418 infection/high intensity.

419 A proportion of pupils in each Primary class in Jaba LGA had urinary  
420 schistosomiasis; however, there were increases in both prevalence and intensity of  
421 the infection from junior classes (Primary 1-3) to senior classes (Primary 4-6).  
422 Though there was no statistical significance in those relationships, the observed  
423 increases could have been due to more water-contact activities among the older  
424 pupils. It shows further that urinary schistosomiasis is unrelated to the class of a  
425 pupil as a risk factor.

426 The highest occurrence of urinary schistosomiasis among pupils in Bitaro, Ankun  
427 and Kwoi could be due to more availability of rivers/streams and the dense water-  
428 contact activities among the pupils compared to other locations. Pupils in Sambang  
429 and Nok recorded an absence of the infection with fewer rivers/streams and reduced  
430 water-contact activities among the pupils; members of these two communities  
431 depend largely on borehole and well water. High water-contact activities, promotes a  
432 higher risk of contracting the infection. However, the highest mean count of *S.*  
433 *haematobium* eggs (of 6.77/10 ml urine) was found in the Central Area and not in the  
434 location of higher occurrence of the infection. Hence, high prevalence of the infection  
435 did not coincide with high intensity of infection because repeated exposure is  
436 required to enhance the development of a heavy infection or high intensity.

437 The overall occurrence of 2.0% heavy infections, 10.3% light infections and egg  
438 count ranging between 0–204eggs/10 ml urine among the pupils in Jaba LGA was  
439 far less than the range of 21-1138eggs/10 ml urine and 62.7% heavy infections  
440 reported by Ossai et al. [26]. This further indicated that pupils in Jaba LGA had  
441 relatively low level of urinary schistosomiasis. Nevertheless, the infected population  
442 continues to pose risk to other uninfected pupils and members of the community.  
443 This will also gradually increase the cost of personal and public health maintenance.

444 The 8.1% prevalence of anaemia among the pupils could have resulted from multiple  
445 sources: dietary deficiency of iron, leukaemia, heavy loss of blood and/or parasitic  
446 diseases. It has been noted that haematological indices of apparently healthy  
447 individuals can be affected by certain factors like age, gender, cultural background,  
448 nature of body build, social activities, nutrition, altitude and other environmental  
449 factors [40]. One of such haematological parameters considered in this study was  
450 PCV and classified as 'anaemic', 'high' or 'normal' [22]. With statistical relevance,  
451 urinary schistosomiasis caused and/or enhanced the development of anaemia  
452 among the pupils. Pupils who become recurrently infected may develop anaemia,  
453 malnutrition and even learning difficulties [41]. The anaemia can be related to the  
454 cumulative loss of blood in terminal haematuria associated with the disease [29] as  
455 well as the continuous feeding on glucose and blood products by schistosomes [38];  
456 making urinary schistosomiasis a chronic symptomatic disease [42].

457 Though schistosomiasis has been reckoned as one of the world's most prevalent  
458 public health problems [37] and Nigeria stands as one of Africa's most endemic  
459 countries [28, 43], with several control/intervention programmes the prevalence of  
460 urinary schistosomiasis seemed not to be decreasing. This is partly due to improper  
461 implementation of the control measures, unawareness and increased water-contact  
462 activities among members of rural communities. In this study, pupils who wash their  
463 clothes in rivers/streams had higher occurrence of the infection than those who wash  
464 at home.

465 The pupils who did not participate in swimming activities were more infected (15.2%)  
466 than those who were swimmers (accounting for 10.7%), but it was not statistically  
467 significant. Some other studies [28], [36] had positively associated schistosomiasis  
468 with water-contact activities (like swimming) in river/stream. The population in this  
469 study might have acquired most the infections through other water-based activities  
470 such as washing or wading in rivers/streams or irrigation farming, indicating that the  
471 disease has multiple means of transmission via skin or body contact with cercariae-  
472 infested water.

473 Pupils' parents' level of formal education did not show any statistical relationship with  
474 urinary schistosomiasis in them. However, the lowest prevalence was observed  
475 among pupils whose parents did not have any formal education. Perhaps, the tight



476 working schedules of educated parents and late return from work gave their  
477 children/wards opportunities to indulge in unsafe water-contact activities. Pupils  
478 whose mothers attained tertiary education had lower infection. However, the  
479 prevalence of the infection increased steadily among the pupils as their fathers' level  
480 of formal education increases. This finding did not agree with the work of Houmsou  
481 et al. [24] who reported that children whose parents did not attain any formal  
482 education as well as those with primary education recorded the highest prevalence  
483 of urinary schistosomiasis.

484 Some symptoms could be useful in the diagnosis of urinary schistosomiasis.  
485 Red/brown-colored urine (a sign of haematuria) and painful micturition witnessed by  
486 some of the subjects in study statistically indicated their associations with the  
487 disease. The association of haematuria with urinary schistosomiasis had been  
488 emphasized by many researchers [36], [44] and described as a classic or main  
489 sign/symptom of the disease [2], [45]. Pains suffered during urination by infected  
490 pupils are direct effects of tissue inflammation, destruction of internal vesical or  
491 ureterovesical sphincter mechanism or minor granulomata due to immunological  
492 reactions to both schistosomes and their eggs [38], [46], [47], [48].

493 The completely zero-level of awareness of schistosomiasis among pupils in Jaba  
494 LGA is indicative of negligence of the disease in Nigeria, a characteristic that defines  
495 it as a 'neglected tropical disease (NTD)' in the condition of poverty [2]. Hence,  
496 unawareness was a constant, and remains a strong risk factor in the continuous  
497 spread of the infection. This opposes the considerable awareness of 74.5% by pupils  
498 in rural communities of Kano State, Nigeria [36] where a lower prevalence of 8.3%  
499 was obtained as against the 12.3% prevalence obtained in this study with complete  
500 unawareness of the infection and its risks. The pupils in Jaba LGA are at continued  
501 risk of getting infected because of unawareness and continuous engagement in  
502 activities in cercariae-infested-water bodies. But it is anticipated that the organized  
503 awareness talks on schistosomiasis in pre-selected schools for this study will help in  
504 reducing the risks of exposure on enlightened pupils.

505

506

507

**508 CONCLUSION**

509 Jaba LGA is considerably an area of mild burden of urinary schistosomiasis,  
510 accounting for prevalence of 12.3%. Female pupils from the LGA had higher  
511 prevalence and intensity of the disease. Hence, they are faced with heightened risks  
512 of developing some complications like female genital schistosomiasis (FGS) and  
513 irreversible infertility.

514 Prevalence and intensity of urinary schistosomiasis had 'wave-like' patterns of  
515 increase with rise in the ages of pupils. Anaemia can result due to urinary  
516 schistosomiasis, a condition which can be severe with high worm burden.

517 One of the basic factors promoting the continuous spread of schistosomiasis is lack  
518 of awareness of the disease, as more children stand at risk in most communities in  
519 Jaba LGA. 'Fadama' farming, painful micturition and red/brown-coloured urine were  
520 statistically associated with urinary schistosomiasis among the study population.

521 In view of the impacts of urinary schistosomiasis on health/development of African  
522 children, concerted efforts by government, school authorities, parents and  
523 researchers should be intensified in the control and prevention of the infection;  
524 treatment of infected/risk groups and early reporting of research/case findings should  
525 be encouraged.

526

**527 CONFLICT OF INTEREST**

528 The authors declare that there is no any financial or conflict of interest.

529

**530 AUTHOR'S CONTRIBUTIONS**

531 Henry Gabriel Bishop

532 Group 1- Concept of the research work, Design of the research work, Acquisition of  
533 data, Data analysis and interpretation

534 Group 2- Drafting of the article manuscript, Critical revision of the article

535 Group 3- Collective approval of final version to published

536

537 Helen Ileigo Inabo

538 Group 1 – Concept of the research work, Design of the research work

539 Group 2 - Critical revision of the article

540 Group 3 - Collective approval of the version to be published

541

542 Ella, Elijah Ekah

543 Group 1 – Concept of the research work, Design of the research work

544 Group 2 - Critical revision of the article

545 Group 3 - Collective approval of the version to be published

546

#### 547 **ACKNOWLEDGEMENTS**

548 We express our appreciation to all the pupils, parents and the management of every  
549 primary school that voluntarily partook in this study. We equally thank Mr. Joseph  
550 Shedow and Mrs. Doris Hassan Dala for their contributions to the success of this  
551 study in Jaba LGA, Kaduna State, Nigeria.

552

#### 553 **REFERENCES**

- 554 1. Osada Y, Kanazawa T. Schistosome: its benefit and harm in suffering from  
555 concomitant diseases. *Journal of Biomedicine and Biotechnology* 2011; ID  
556 264173: 1-10.
- 557 2. World Health Organisation. Schistosomiasis Fact Sheet Updated February  
558 2016. Assessed on 1<sup>st</sup> April 2016, from:  
559 <http://www.who.int/mediacentre/factsheets/fs115/en/>
- 560 3. Edward JP, Andrew SW. Immunology of schistosomiasis. *Nature Reviews*  
561 *Immunology* 2002; 2:499-511.
- 562 4. Cox FEG. *Modern Parasitology: a Text Book of Parasitology*, 2<sup>nd</sup> ed.  
563 Germany: Backwell Science Limited, 2004.
- 564 5. Noble ER, Glem AN. *Biology of Animal Parasites*, 5<sup>th</sup> ed. Philadelphia, USA:  
565 Lea and Febiger: 1982. P.157-158.
- 566 6. Ibironke O, Koukounari A, Asaolu S, Moustaki I, Shiff C. Validation of a new  
567 test for *Schistosoma haematobium* based on detection of Dra 1 DNA  
568 fragments in urine: Evaluation through latent class analysis. *PloS Neglected*  
569 *Tropical Diseases* 2012; 6(1): e1464.

- 570 7. Kanwai S, Ndams IS, Kogi E, Gyem ZG, Hena JS. Urinary schistosomiasis  
571 infection in Dumbin Dustse, Igabi Local Government Area, Kaduna State,  
572 Nigeria. *Science World Journal* 2011; 6(3):1-3.
- 573 8. Elele K, Ewurum N. Current Status of Urinary Schistosomiasis in Four  
574 Communities (Obedum, Anyu, Amerikpoko, and Odau) In Abual/Odual, Rivers  
575 State, South-South, Nigeria. *Reiko International Journal of Science and*  
576 *Technology* 2013; 4(2): Assessed from: [http://reikojournals.org/index.php/the-](http://reikojournals.org/index.php/the-journal-of-social-and-economic-3?id=214)  
577 [journal-of-social-and-economic-3?id=214](http://reikojournals.org/index.php/the-journal-of-social-and-economic-3?id=214)
- 578 9. Rollinson DA, Southgate VR. The genus *Schistosoma*: a taxonomic appraisal.  
579 In Rollinson D, Simpson AJ. (eds). *The Biology of Schistosome: from Genes*  
580 *to Latrines*. London: Academic Press: 1987. p.1-49.
- 581 10. Jamjoom MB. Molecular identification of some *Schistosoma mansoni* isolates  
582 in Saudi Arabia. *World Journal of Medical Sciences* 2006; 1(2):102-107.
- 583 11. World Health Organisation. Schistosomiasis Fact sheet N°115, Updated May  
584 2015. Assessed on 27<sup>th</sup> June 2016 from:  
585 <http://www.who.int/mediacentre/factsheets/fs115/en/>
- 586 12. Clerinx J, Soentjens P. Epidemiology, pathogenesis, and clinical  
587 manifestations of schistosomiasis.2015. Assessed on 27<sup>th</sup> June 2016 from:  
588 [http://www.uptodate.com/contents/epidemiology-pathogenesis-and-clinical-](http://www.uptodate.com/contents/epidemiology-pathogenesis-and-clinical-manifestations-of-schistosomiasis)  
589 [manifestations-of-schistosomiasis](http://www.uptodate.com/contents/epidemiology-pathogenesis-and-clinical-manifestations-of-schistosomiasis)
- 590 13. Barsoum RS. Urinary schistosomiasis review. *Journal of Advanced Research*  
591 2013; 4:453-459.
- 592 14. Sakanari JA, Mckerrow JH. Medical Parasitology. In Brooks GF, Carroll KC,  
593 Butel JS, Morse SA, Mietzner TA. (eds). *Jawertz, Melnick and Adelberg's*  
594 *Medical Microbiology*, 25<sup>th</sup> ed. US: McGraw Hill Companies, Inc., 2010. p.665-  
595 701.
- 596 15. Rollinson DA. Wake up call for urinary schistosomiasis: reconciling research  
597 effort with public health importance. *Parasitology* 2009; 136:1593-1610.
- 598 16. Rinaldi G, Okatcha TI, Popratiloff A, Ayuk MA, Suttiprapa S, Mann VH, Liang  
599 Y, Lewis FA, Loukas A, Brindley PJ. Genetic manipulation of *Schistosoma*  
600 *haematobium*, the neglected schistosome. *PloS Neglected Tropical Diseases*  
601 2011; 5:e1348.

- 602 17. Brindley PJ, Hotez PJ. Break out: urogenital schistosomiasis and  
603 Schistosoma haematobium infection in the post-genomic era. PloS Neglected  
604 Tropical Diseases 2013, 7(3):e1961.
- 605 18. Chidozie EU, Duniyan SY. Urinary schistosomiasis epidemiological survey of  
606 urinary schistosomiasis among children in selected schools: A preliminary  
607 study in Minna, Nigeria. African Journal of Biotechnology 2008; 7(16):2773-  
608 2776.
- 609 19. World Health Organisation. Report of the WHO informal consultation on  
610 Schistosomiasis Control. Geneva, 2-4 December, 1998. Geneva:  
611 WHO/CDS/CPC/SIP/99.2. 1999. Assessed from:  
612 [http://apps.who.int/iris/bitstream/10665/65978/1/WHO\\_CDS\\_CPC\\_SIP\\_99.2.p](http://apps.who.int/iris/bitstream/10665/65978/1/WHO_CDS_CPC_SIP_99.2.pdf)  
613 [df](http://apps.who.int/iris/bitstream/10665/65978/1/WHO_CDS_CPC_SIP_99.2.pdf)
- 614 20. Uneke C, Ugwuoke-Adibuah S, Nwakpu K, Ngwu B. An Assessment of  
615 Schistosoma haematobium infection and urinary tract bacterial infection  
616 among school children in rural eastern Nigeria. The Internet Journal of  
617 Laboratory Medicine 2009; 4(1):1-6.
- 618 21. Nokculture.com (n.d). Nok Culture. Assessed on 4<sup>th</sup> December, 2015 from:  
619 <http://nokculture.com/>
- 620 22. Cheesbrough M. District Laboratory Practice in Tropical Countries, Part I, 2<sup>nd</sup>  
621 ed. updated. Cambridge, UK: Cambridge University Press: 2009.
- 622 23. Ifeanyi CIC, Matur BM, Ikeneche NF. Urinary schistosomiasis and  
623 concomitant bacteriuria in the Federal Capital Territory Abuja Nigeria. New  
624 York Science Journal 2009; 2(2):1-8.
- 625 24. Houmsou RS, Amuta EU Sar TT. Profile of an epidemiological study of urinary  
626 schistosomiasis in two local government areas of Benue State, Nigeria.  
627 International Journal of Medicine and Biomedical Research 2012; 1(1):39-48.
- 628 25. Oluwasogo OA, Fagbemi OB. Prevalence and risk factors of Schistosoma  
629 haematobium infections among primary school children in Igbokuta Village,  
630 Ikorodu North Local Government, Lagos State. IOSR Journal of Nursing and  
631 Health Science 2013; 2(6): 62-68.
- 632 26. Ossai OP, Dankoli R, Nwodo C, Tukur D, Nsubuga P, Ogbuabor D, Ekwueme  
633 O, Abonyi G, Ezeanolue E, Nguku P, Nwagbo D, Idris S, Eze G. Bacteriuria

- 634 and urinary schistosomiasis in primary school children in rural communities in  
635 Enugu State, Nigeria, 2012. *The Pan African Medical Journal* 2014; 18(1):15.
- 636 27. Balla HJ, Babaganal Baba S, Ibrahim H. Incidence of urinary schistosomiasis  
637 among out-of-school pupils and “Almajiris” in Dikwa, North Eastern Nigeria.  
638 *Global Journal of Medical Research (C)* 2015; 15(2):9-13.
- 639 28. Omenesa HO, Bishop HG. Raji HM. Prevalence of Urinary Schistosomiasis  
640 among Pupils Attending Primary Schools in Bomo Village, Zaria-Nigeria.  
641 *International Journal of Research in Engineering and Science* 2015; 3(5):14-  
642 19.
- 643 29. Grimes JET, Croll D, Harrison WE, Utzinger J, Freeman MC, Templeto MR.  
644 The Relationship between Water, Sanitation and Schistosomiasis: A  
645 Systematic Review and Meta-analysis. *PLOS Neglected Tropical Diseases*  
646 2014; 8(12): e3296. doi: 10.1371/journal.pntd.0003296.
- 647 30. Dabo A, Badawi HM, Bary B, Doumbo OK. Urinary schistosomiasis among  
648 preschool-aged children in Sahelian rural communities in Mali. *Parasite and*  
649 *Vectors* 2011; 4(21):1-7.
- 650 31. Tay SCK, Amankwa R, Gbedema SY. Prevalence of *Schistosoma*  
651 *haematobium* infection in Ghana: a retrospective case study in Kumasi.  
652 *International Journal of Parasitology Research* 2011; 3(2):48-52.
- 653 32. Tetteh-Quarcoo PB, Attah SK, Donkor ES, Nyako M, Minamor AA, Afutu E,  
654 Hervie ET, Ayeh-Kumi PF. Urinary schistosomiasis in children—Still a  
655 concern in part of the Ghanaian capital city. *Open Journal of Medical*  
656 *Microbiology* 2013; 3:151-158.
- 657 33. Geleta S, Alemu A, Getie S, Mekonnen Z Erk B. Prevalence of urinary  
658 schistosomiasis and associated risk factors among Abobo primary school  
659 children in Gambella Regional State, Southwestern Ethiopia: a cross sectional  
660 study. *Parasites & Vectors* 2015; 8(215):1-9. doi 10.1186/s13071-015-0822-5.
- 661 34. Samie A, Nchachi DJ, Obi CL, Igumbor EO. Prevalence and temporal  
662 distribution of *Schistosoma haematobium* infections in the Vhembe district,  
663 Limpopo Province, South Africa. *African Journal of Biotechnology* 2010;  
664 9(42):7157- 7164. doi: 10.5897/AJB10.015.

- 665 35. Ingang-Etoh PC, Essien UC, Amama SA, Useh MF. Prevalence of urinary  
666 schistosomiasis among school children in Ukwelo-Obudu and Abini  
667 communities in Cross River State, Nigeria. *Port Harcourt Medical Journal*  
668 2009; 3(3): assessed on 14<sup>th</sup> January, 2016 from:  
669 <http://www.ajol.info/index.php/phmedj/article/view/45251>
- 670 36. Dawaki S, Al-Mekhlafi HM, Ithoi I, Ibrahim J, Abdulsalam AM, Ahmed, A, Sady  
671 H, Nasr NA, Atroosh WM. The Menace of schistosomiasis in Nigeria:  
672 knowledge, attitude, and practices regarding schistosomiasis among rural  
673 communities in Kano State. *PLoS ONE* 2015; 10(11): e0143667.  
674 doi:10.1371/journal.pone.0143667.
- 675 37. Bigwan EI, Kunihya, RZ, John TJ. Epidemiological survey of urinary  
676 schistosomiasis among primary school children in Michika, Adamawa State,  
677 North-Eastern Nigeria. *International Journal of Current Research and Review*  
678 2013; 5(5):111-116.
- 679 38. Leder K, Weller PF. Epidemiology, pathogenesis and clinical features of  
680 schistosomiasis 2011. Assessed on 3<sup>rd</sup> November, 2015 from:  
681 <http://cursoenarm.net/UPTODATE/contents/mobipreview.htm?37/42/38560?s>  
682 [ource=see\\_link](#)
- 683 39. Kjetland EF, Leutscher PD, Ndhlovu PD A review of female schistosomiasis.  
684 *Trends in Parasitology* 2012; 28(2):58-65. doi: 10.1016/j.pt.2011.10.008.
- 685 40. El-Hazmi MAF, Warsy AS. Normal reference values for hematological  
686 parameters, red cell indices, HB A2 and HB F from early childhood through  
687 adolescence in saudis. *Annals of Saudi Medicine* 2001; 21(3-4):165-169.
- 688 41. Centers for Disease Control and Prevention (CDC). Parasite-Schistosomiasis.  
689 2012. Assessed on 8<sup>th</sup> January, 2016 from  
690 <http://www.cdc.gov/parasites/schistosomiasis/biology.html>.
- 691 42. King CH. It's time to Dispel the Myth of "Asymptomatic schistosomiasis. *PloS*  
692 *Neglected Tropical Diseases* 2015; 9(2):e0003504.
- 693 43. Chitsulo L, Engels D, Montresor A, Savioli L. The global status of  
694 schistosomiasis and its control. *Acta Tropica* 2000; 77:41-51.
- 695 44. Morenikeji O, Quazim J, Omoregie C, Hassan A, Nwuba R, Anumudu C,  
696 Adejuwon S, Salawu O, Jegede A, Odaibo, A. A cross-sectional study on

- 697 urogenital schistosomiasis in children; haematuria and proteinuria as  
698 diagnostic indicators in an endemic rural area of Nigeria. *African Health*  
699 *Science* 2014; 14(2):390–396.
- 700 45. Van der Werf MJ, de Vlas SJ. Diagnosis of urinary schistosomiasis: a novel  
701 approach to compare bladder pathology measured by ultrasound and three  
702 methods for hematuria detection. *American Journal of Tropical Medicine and*  
703 *Hygiene* 2004; 7(1):98-106.
- 704 46. Elbaz T, Esmat G. Hepatic and intestinal schistosomiasis: review. *Journal of*  
705 *Advanced Research* 2013; 4:445-452.
- 706 47. Botelho MC, Figueiredo J, Alves H. Bladder cancer and urinary  
707 schistosomiasis in Angola. *Journal of Nephrology Research* 2015; 1(1):22-24.  
708 doi:10.17554/j.issn.2410-0579.2015.01.4.
- 709 48. Botelho MC, Machado JC, Brindley PJ, Correia da Costa JM. Targeting  
710 molecular signaling pathways of *Schistosoma haematobium* infection in  
711 bladder cancer. *Virulence* 2011; 2:267-279.
- 712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728



729 **TABLES**

730

731 Table 1: Occurrence and Intensity of Urinary Schistosomiasis according to Class of  
732 Pupils

733

<b>Primary Class</b>	<b>No. of Samples Examined</b>	<b>* Positive Cases No (%)</b>	<b>Range of Count</b>	<b>Female Mean of Eggs/10ml urine</b>	<b>Male Mean of Eggs/10ml urine</b>	<b>Class Mean of Eggs/10ml urine</b>
<b>1</b>	74	7(9.5)	0-5	0.34	0.14	0.24
<b>2</b>	79	5(6.3)	0-103	3.32	0.56	1.65
<b>3</b>	84	8(9.5)	0-62	2.48	0.19	1.50
<b>4</b>	104	15(14.4)	0-130	4.13	2.91	3.47
<b>5</b>	92	15(16.3)	0-78	5.14	2.09	3.72
<b>6</b>	72	12(16.7)	0-204	9.74	0.38	5.32
<b>Total</b>	<b>505</b>	<b>62(12.3)</b>	<b>0-204</b>	<b>4.18</b>	<b>1.22</b>	<b>2.69</b>

734

735 \* $\chi^2 = 6.850$ ; df = 5; P = 0.232, LR = 7.178; df = 5 P = 0.208

736

737

738

739

740 Table 2: Gender-related Distribution of Urinary Schistosomiasis among Pupils in  
741 Jaba LGA

742

Gender	No. of Samples Examined	No. Positive (%)	No. Negative (%)
Female	251	39 (15.5)	212 (84.5)
Male	254	23 (9.1)	231 (90.9)
<b>Total</b>	<b>505</b>	<b>62 (12.3)</b>	<b>443 (87.7)</b>

743

744  $\chi^2 = 4.926$ ;  $df = 1$ ;  $P = 0.026$ ,  $LR = 4.973$ ;  $df = 1$ ;  $P = 0.026$

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765 Table 3: Occurrence and Mean Intensity of Urinary Schistosomiasis according to  
766 Sampling Locations

767

<b>Sampling Location</b>	<b>No. of Samples examined</b>	<b>*Prevalence No. (%)</b>	<b>**Intensity (Mean of Eggs/10ml urine)</b>	<b>Standard Error of Mean</b>
<b>Ankun</b>	27	6 (22.2)	3.30	2.301
<b>Bitaro</b>	112	26(23.2)	4.85	2.018
<b>Central Area</b>	57	7(12.3)	6.77	3.335
<b>Chori</b>	30	1(3.3)	0.10	0.100
<b>Dura</b>	21	1(4.8)	0.29	0.286
<b>Gora</b>	26	4(15.4)	0.73	0.358
<b>Kwoi</b>	59	12(20.3)	2.73	1.451
<b>Nok</b>	33	0(0.0)	0.00	0.000
<b>Sambang</b>	78	0(0.0)	0.00	0.000
<b>Yadi-Pyok</b>	62	5(8.1)	2.47	1.443

768

769 \*  $\chi^2 = 38.599$ ;  $df = 9$ ;  $p = 0.000$ ,  $LR = 50.200$ ;  $df = 9$ ;  $p = 0.000$

770 \*\*ANOVA Value (F) = 1.419;  $df = 9$ ;  $P = 0.177$

771

772

773

774

775

776

777

778

779

780

781 Table 4: Distribution of level of urinary schistosomiasis across sampling locations in  
 782 Jaba LGA, Kaduna State, Nigeria  
 783

Level of Infection	Samples Size	Ankun	Bitaro	Central Area	Chori	Dura	Gora	Kwoi	Nok	Sambang	Yadi-Pyok
None	443	21 (4.7)	86 (19.4)	50 (11.3)	29 (6.5)	20 (4.5)	22 (5.0)	47 (10.6)	33 (7.4)	78 (17.6)	57 (12.9)
Light	52	5 (9.6)	23 (44.2)	3 (5.8)	1 (1.9)	1 (1.9)	4 (7.7)	11 (21.2)	0 (0.0)	0 (0.0)	4 (7.7)
Heavy	10	1 (10.0)	3 (30.0)	4 (40.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (10.0)	0 (0.0)	0 (0.0)	1 (10.0)
<b>Total</b>	<b>505</b>	<b>27</b> <b>(5.3)</b>	<b>112</b> <b>(22.2)</b>	<b>57</b> <b>(11.3)</b>	<b>30</b> <b>(5.9)</b>	<b>21</b> <b>(4.2)</b>	<b>26</b> <b>(5.1)</b>	<b>59</b> <b>(11.7)</b>	<b>33</b> <b>6.5)</b>	<b>78</b> <b>(15.4)</b>	<b>62</b> <b>(12.3)</b>

784  
 785  
 786  
 787  
 788  
 789  
 790  
 791

$\chi^2 = 50.094$ , df = 18, P = 0.000, LR = 59.532; df = 18; P = 0.000

792 Table 5: Age-related distribution of the level of urinary schistosomiasis among pupils  
 793 in Jaba LGA, Kaduna State, Nigeria

794

Age (Years)	No. of Samples Examined	No Infection No. (%)	Light Infection No. (%)	Heavy Infection No. (%)
4	2	2 (100.0)	0 (0.0)	0 (0.0)
5	10	10 (100.0)	0 (0.0)	0 (0.0)
6	19	19 (100.0)	0 (0.0)	0 (0.0)
7	43	40 (93.0)	2 (4.7)	1 (2.3)
8	67	64 (95.5)	3 (4.5)	0 (0.0)
9	75	65 (86.7)	9 (12.0)	1 (1.3)
10	97	81 (83.5)	13 (13.4)	3 (3.1)
11	77	71 (92.2)	6 (7.8)	0 (0.0)
12	54	47 (87.0)	6 (11.1)	1 (1.9)
13	34	25 (73.5)	6 (17.6)	3 (8.8)
14	19	16 (84.2)	3 (15.8)	0 (0.0)
15	8	3 (37.5)	4 (50.0)	1 (12.5)
<b>Total</b>	<b>505</b>	<b>443 (87.7)</b>	<b>52 (10.3)</b>	<b>10 (2.0)</b>

795

796  $\chi^2 = 44.715$  df = 22; P = 0.003, LR = 40.665 df = 22; P = 0.009

797

798

799

800

801

802 Table 6: Relationship between urinary schistosomiasis and anaemia among pupils  
 803 in Jaba LGA

804

<b>Status of Urinary Schistosomiasis</b>	<b>No. of Samples Examined</b>	<b>Anaemic PCV No. Positive (%)</b>	<b>Normal PCV No. Positive (%)</b>	<b>High PCV No. Positive (%)</b>
<b>Not Infected</b>	443	30 (6.8)	163 (36.8)	250 (56.4)
<b>Infected</b>	62	11 (17.7)	27 (43.5)	11 (38.7)
<b>Total</b>	<b>505</b>	<b>41 (8.1)</b>	<b>190 (37.6)</b>	<b>274 (54.3)</b>

805

806  $\chi^2 = 11.870$ ; df = 2; P = 0.003, LR = 10.401; df = 2; P = 0.006

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825 Table 7: Effect of the level of urinary schistosomiasis on pupils' packed cell volume

826

Level of Infection	No. of Samples Examined	Anaemic PCV No. Positive (%)	Normal PCV No. Positive (%)	High PCV No. Positive (%)	Total
<b>None</b>	443	30 (6.8)	163 (36.8)	250 (56.4)	<b>443 (87.7)</b>
<b>Light</b>	52	9 (17.3)	24 (46.2)	19 (36.5)	<b>52 (10.3)</b>
<b>Heavy</b>	10	2 (20.0)	3 (30.0)	5 (50.0)	<b>10 (2.0)</b>
<b>Total</b>	<b>505</b>	<b>41(8.1)</b>	<b>190 (37.6)</b>	<b>274 (54.3)</b>	<b>505 (100.0)</b>

827

828  $\chi^2 = 12.807$ ; df = 4; P = 0.012, LR = 11.353; df = 4; P = 0.023

829

830

831

832

833

834

835

836

837

838

839

840

841

842 Table 8: Relationship between the level of urinary schistosomiasis and gender of  
 843 pupils

844

<b>Level of Infection</b>	<b>No. of Samples Examined</b>	<b>Female No. Positive (%)</b>	<b>Male No. Positive (%)</b>	<b>Total</b>
<b>None</b>	443	212 (47.9)	231 (52.1)	<b>443 (87.7)</b>
<b>Light</b>	52	32 (61.5)	20 (38.5)	<b>52 (10.3)</b>
<b>Heavy</b>	10	7 (70.0)	3 (30.0)	<b>10 (2.0)</b>

845

846  $\chi^2 = 5.166$ ;  $df = 2$ ;  $P = 0.076$ ,  $LR = 5.237$ ;  $df = 0.073$

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863



864 Table 9: Effect of level of urinary schistosomiasis on occurrence of haematuria  
 865 among pupils in Jaba LGA, Kaduna State

866

Level of infection	No. of Samples Examined	Absence of Haematuria No. Positive (%)	Presence of Haematuria No. Positive (%)
None	443	423 (95.5)	20 (4.5)
Light	52	47 (90.4)	5 (9.6)
Heavy	10	10 (100.0)	0 (0.0)
<b>Total</b>	<b>505</b>	<b>480 (95.0)</b>	<b>25 (5.0)</b>

867

868  $\chi^2 = 3.105$ ; df = 2; P = 0.212, LR = 3.108; df = 2; P = 0.211

869

870

871

872

873

874

875

876

877

878

879

880

881

882 Table 10: Risk Factors associated with Urinary Schistosomiasis among Primary  
883 School Pupils in Jaba LGA

884

Risk Factor	Response Category	No. of Samples Examined	Positive No. (%)	Negative No. (%)
<b>Awareness of Schistosomiasis<sup>#</sup></b>	Unaware	505	62 (12.3)	443 (87.7)
	Aware	0	0 (0.0)	0(0.0)
<b>'Fadama' Farming<sup>*</sup></b>	No	376	34 (9.0)	342 (91.0)
	Yes	129	28 (21.7)	101 (78.3)
<b>Fishing in River/Stream<sup>**</sup></b>	No	396	52 (13.1)	344 (86.9)
	Yes	109	10 (9.2)	99 (90.8)
<b>Swimming in River/Stream<sup>†</sup></b>	No	178	27 (15.2)	151 (84.8)
	Yes	327	35 (10.7)	292 (89.3)
<b>Place of Laundering<sup>@</sup></b>	Home	315	34 (10.8)	281 (89.2)
	River/Stream	190	28 (14.7)	162 (85.3)

885

886 <sup>#</sup>No computed statistics because unawareness was a constant.

887 <sup>\*</sup>  $\chi^2 = 14.300$ ; df = 1; P = 0.000, LR = 12.915; df = 1; P = 0.000

888 <sup>\*\*</sup>  $\chi^2 = 1.243$ ; df = 1; P = 0.265, LR = 1.319; df = 1; P = 0.251

889 <sup>†</sup>  $\chi^2 = 2.134$ ; df = 1; P = 0.144, LR = 2.080; df = 1; P = 0.149

890 <sup>@</sup>  $\chi^2 = 1.711$ ; df = 1; P = 0.191, LR = 1.679; df = 1; P = 0.195

891

892

893

894 Table 11: Association of parental level of formal education and urinary  
 895 schistosomiasis status of pupils in Jaba LGA

896

Parent Category	Level of Formal Education	No. of Samples Examined	No. Positive (%)	No. Negative (%)
<b>Mother*</b>	None	58	2 (3.4)	56 (96.6)
	Primary	94	13 (13.8)	81 (86.2)
	Secondary	302	43 (14.2)	259 (85.8)
	Tertiary	51	4 (7.8)	47 (92.2)
	<b>Total</b>	<b>505</b>	<b>62 (12.3)</b>	<b>443 (87.7)</b>
<b>Father**</b>	None	59	4 (6.8)	55 (93.2)
	Primary	63	6 (9.5)	57 (90.5)
	Secondary	299	40 (13.4)	259 (86.6)
	Tertiary	84	12 (14.3)	72 (85.7)
	<b>Total</b>	<b>505</b>	<b>62 (12.3)</b>	<b>443 (87.7)</b>

897

898 \*  $\chi^2 = 6.418$ ; df = 3; P = 0.093, LR = 7.947; df = 3; P = 0.047

899 \*\*  $\chi^2 = 2.750$ ; df = 3; P = 0.432, LR = 3.041; df = 3; P = 0.385

900

901

902

903 Table 12: Sign/symptoms associated with urinary schistosomiasis among the pupils  
 904 in Jaba LGA, Kaduna State, Nigeria.

905

Symptom	Category	No. of Samples Examined	No. (%) Positive	No. (%) Negative
<b>Haematuria*</b>	No	480	57 (11.9)	423 (88.1)
	Yes	25	5 (20.0)	20 (80.0)
<b>Painful micturition**</b>	No	473	54 (11.4)	419 (88.6)
	Yes	32	8 (25.0)	24 (75.0)
<b>Frequent micturition<sup>a</sup></b>	No	496	61 (12.3)	435 (87.7)
	Yes	9	1 (11.1)	8 (88.9)
<b>Abdominal pain<sup>a</sup></b>	No	496	61 (12.3)	435 (87.7)
	Yes	9	1 (11.1)	8 (88.9)
<b>Fever<sup>b</sup></b>	No	487	58 (11.9)	429 (88.1)
	Yes	18	4 (22.2)	14 (77/8)
<b>Urine colour<sup>c</sup></b>	Brown and cloudy	2	1 (50.0)	1 (50.0)
	Cloudy	39	5 (12.8)	34 (87.2)
	Milky-white	91	9 (9.9)	82 (90.1)
	Red and cloudy	92	23 (25.0)	69 (75.0)
	Yellow-orange	281	24 (8.5)	257 (91.5)

906

907 \*  $\chi^2 = 1.457$ ; df = 1; P = 0.227, LR = 1.266; df = 1; P = 0.260

908 \*\*  $\chi^2 = 5.135$ ; df = 1; P = 0.023, LR 3.951; df = 1; P = 0.041

909 <sup>a</sup>  $\chi^2 = 0.012$ ; df = 1; P = 0.914, LR = 0.012, df = 1; P = 0.913

910 <sup>b</sup>  $\chi^2 = 1.714$ ; df = 1; P = 0.190, LR = 1.439; df = 1; P = 0.230

911 <sup>c</sup>  $\chi^2 = 20.604$ ; df = 4; P = 0.000, LR = 17.315; df = 4; P = 0.02

912

913 **FIGURE LEGENDS**

914

915 Figure 1: Overall prevalence of urinary schistosomiasis among pupils of Jaba LGA of  
916 Kaduna State, Nigeria.

917

918 Figure 2: Age prevalence of urinary schistosomiasis among pupils in Jaba LGA

919

920 Figure 3: Age pattern of intensity of urinary schistosomiasis

921

922 Figure 4: Prevalence of anaemia among the pupils in Jaba LGA, Kaduna State,  
923 Nigeria.

924

925 Figure 5: Various appearances of *S. haematobium* ova in urine samples of pupils  
926 from Jaba LGA, Kaduna State, Nigeria. (A, and B) wet mounts stained with Lugol's  
927 iodine; (C and D) - without Lugol's iodine on 40x objective)

928

929 Figure 5: Various appearances of *S. haematobium* ova in urine samples of pupils  
930 from Jaba LGA, Kaduna State, Nigeria. (A, and B) wet mounts stained with Lugol's  
931 iodine; (C and D) - without Lugol's iodine on 40x objective)

932

933

934

935

936

937

938

939

940

941


942

943

944


945 **FIGURES**

946



947

948 Figure 1: Overall prevalence of urinary schistosomiasis among pupils of Jaba LGA of  
949 Kaduna State, Nigeria.




950  
951

952 ( $t$ -test = 4.251, df = 503,  $p$  = 0.000)

953 Figure 2: Age prevalence of urinary schistosomiasis among pupils in Jaba LGA

954




955  
956

957 Spearman's correlation coefficient ( $r_s$ ) = 0.172, P = 0.000; Pearson correlation  
958 coefficient, ( $r$ ) = 0.131, P = 0.003

959 Figure 3: Age pattern of intensity of urinary schistosomiasis






960  
961

962 Figure 4: Prevalence of anaemia among the pupils in Jaba LGA, Kaduna State,  
963 Nigeria.

964 **Keys:** Anaemic PCV was < 34%, Normal PCV was  $\geq 34 \leq 45\%$ , High was  $\geq 46\%$



965

966

967 Figure 5: Various appearances of *S. haematobium* ova in urine samples of pupils  
968 from Jaba LGA, Kaduna State, Nigeria. (A, and B) wet mounts stained with Lugol's  
969 iodine; (C and D) - without Lugol's iodine on 40x objective)

970

971

972

973



974

975

976 Figure 6: River Dhugk Gira and River Langk Wada, the most popular rivers in Bitaro,  
977 Jaba LGA, (notably for swimming, fishing and washing by pupils and the members of  
978 the community).